

ASTHMA ACTION PLAN

Photo

Student Information:		
Student:	Birthdate:	
School:	Grade/Rm	
Emergency Contact Information:		
Name	Relationship	Telephone number
1		
2		
3		
Healthcare Provider	Tel	

Asthma Emergency Action:

The following are possible signs of an asthma emergency:

- Difficulty breathing, walking, or talking
- Blue or gray discoloration of the lips or fingernails
- Failure of medication to reduce worsening symptoms.

These signs indicate the need for emergency medical care. The steps that should be taken:

- Activate the emergency medical system in your area. Call 911.
- Call Parent/Guardian and/or Healthcare Provider •

Triggers:

Name of Medication	Dosage	Time

Start Date End Date

Steps for an Acute Asthma Episode (to be completed by prescriber)

- 1. Administer Inhaler as ordered.
- 2. If no relief in _____ minutes, Repeat Inhaler as ordered.
- 3. Contact Parent/Emergency Contact
- 4. Call 911
- 5. Additional Instructions:

Please check if STUDENT is permitted by healthcare provider to CARRY an inhaler and SELF-MEDICATE at school. (See next page.) (In accordance with ORC 3313.716/3313.14)

Signature of Prescriber	Date	
Signature of Parent/Guardian_	 Date	

Adverse reactions that should be reported to physician:

- Chest pain.
- Rash, hives, or itching.
- Fast, pounding, or irregular heartbeat.
- Swelling of the face, throat, tongue, lips, eyes, hands, feet, ankles, or lower legs.
- Difficulty swallowing.
- Worsened breathing.
- Hoarseness.

Adverse reactions for unauthorized user:

- Racing heart beat
- Feeling very shaky

In the event that medication does not produce the expected relief from student's asthma attack, follow the "Steps for an Acute Asthma Episode" (on first page)

Other special instructions:

Copies must be provided to the principal and to the nurse.

Reviewed by Dr. Carly Wilbur April 2019

